
Name 10 features associated, directly or indirectly, with liver disease which can be
found in the hand. 10
Anaemia
Jaundice
Bruising
Palmar erythaema
Dupitrien's contracture
Leuconychia
Tremor
Tattoos
Clubbing
Spider naevi



Junctional Tachycardia (½ point for supraventricular), Rate of 115 (110-120), Left Axis Deviation, retrograde p waves buried within QRS

According to the APLS SVT algorithm what is the first treatment in children with SVT? 1 point Which 6 drugs can be used to treat SVT? 6 points

Vagal manoeuvre
Adenosine50/100/250 mcg/kg
Verapamil
Digoxin
Porpranolol 50mcg/kg
Flecanide 2mg/kg
Amiodarone 5mg/kg

EXTRA ½ POINTS FOR THE DOSES

Which of the above drugs should be avoided in children under 1?

Verapamil

A 3 year old boy present to your department because his parents noticed him limping over the last 3 days. There is no history of trauma and he had been walking and running around normally until now. Name 10 acquired, non traumatic, causes of a limp in a child which are localised to the leg- 5 points
Cellulitis
Septic arthritis
Osteomyelitis
Perthe's
Irritable hip
Juvenile RA
Neoplasms
Warts Rickets
Scurvy
Polio
A 2 year old boy presents to A&E with 1 day H?O NWB on his L foot. There is a vague history of possible fall
yesterday at grandparents but he was walking well all day until developing a slight limp in the evening. He has
chicken pox all over his body, temperature of 38.4 °C, tachycardia of 140 regular. He is covered in ch. Pox rash
and slightly flushed on his face. He is refusing to put any wt on his L leg. On close examination he appears to
be tender just above his lateral malleolus with a slight swelling there. His hip, knee etc appear to be fine.
List 3 investigations which are most likely to give you the diagnosis? 3 points X-ray, USS, bone scan, ½ point for MRI or CT
A-ruy, Oss, vone scan, 72 point for WICI or CI
Give 2 most likely diagnosis 2 points Fractive distal fibrila, estacomolitis fibrila, there is an association between ab new and estacomolitis/sentia
Fracture distal fibula, osteomyelitis fibula- there is an association between ch pox and osteomyelitis/septic arthritis